

Literaturliste zum Artikel:

## Kieferorthopädie als Fachdisziplin im Zentrum für Seltene Erkrankungen

Dr. Teresa Kruse, Prof. Dr. Bert Braumann

- [1] Melnikova I. Rare diseases and orphan drugs. *Nature Reviews Drug Discovery* 2012; 11: 267
- [2] [https://www.bfarm.de/DE/Kodiersysteme/Kooperationen-und-Projekte/Orphanet-Deutschland/ node.html](https://www.bfarm.de/DE/Kodiersysteme/Kooperationen-und-Projekte/Orphanet-Deutschland/node.html) last accessed: 03/30/2022
- [3] <https://www.sozialministerium.at/Themen/Gesundheit/Seltene-Krankheiten.html> last accessed: 03/30/2022
- [4] <http://www.orpha.net/national/CH-DE/index/startseite/> last accessed: 03/30/2022
- [5] <http://www.nrw-zse.de/> last accessed: 03/30/2022
- [6] <https://ern-cranio.eu/> last accessed: 03/30/2022
- [7] Cobourne MT. The complex genetics of cleft lip and palate. *Eur J Orthod* 2004; 26: 7-16. doi:10.1093/ejo/26.1.7
- [8] Mossey PA, Little J, Munger RG et al. Cleft lip and palate. *The Lancet* 2009; 374: 1773-1785
- [9] Mangold E, Ludwig KU, Nothen MM. Breakthroughs in the genetics of orofacial clefting. *Trends Mol Med* 2011; 17: 725–733. doi:10.1016/j.molmed.2011.07.007
- [10] Stuppia L, Capogreco M, Marzo G et al. Genetics of syndromic and nonsyndromic cleft lip and palate. *Journal of Craniofacial Surgery* 2011; 22: 1722-1726
- [11] Scott AR, Mader NS. Regional variations in the presentation and surgical management of Pierre Robin sequence. *Laryngoscope* 2014; 124: 2818-2825. doi:10.1002/lary.24782
- [12] Vatlach S, Maas C, Poets CF. Birth prevalence and initial treatment of Robin sequence in Germany: a prospective epidemiologic Study. *Orphanet J Rare Dis* 2014; 9:9. doi:10.1186/1750-1172-9-9
- [13] Evans AK, Rahbar R, Rogers GF et al. Robin sequence: a retrospective review of 115 patients. *Int J Pediatr Otorhinolaryngol* 2006; 70: 973-980. doi:10.1016/j.ijporl.2005.10.016
- [14] Breugem CC, Evans KN, Poets CF et al. Best Practices for the Diagnosis and Evaluation of Infants With Robin Sequence: A Clinical Consensus Report. *JAMA Pediatr* 2016; 170: 894-902. doi:10.1001/jamapediatrics.2016.0796

- [15] Evans KN, Sie KC, Hopper RA et al. Robin sequence: from diagnosis to development of an effective management plan. *Pediatrics* 2011; 127: 936-948. doi:10.1542/peds.2010-2615
- [16] Robin P. Glossoptosis due to atresia and hypotrophy of the mandible. *American journal of diseases of children* 1934; 48: 541—547
- [17] Reddy VS. Evaluation of upper airway obstruction in infants With Pierre Robin sequence and the role of polysomnography Review of current evidence. *Paediatr Respir Rev* 2016; 17: 80—87. doi:10.1016/j.prrv.2015.10.001
- [18] Neuschulz J, Wilhelm L, Christ H et al. Prenatal indices for mandibular retrognathia/micrognathia. *J Orofac Orthop* 2015; 76: 30—40. doi:10.1007/s00056-014-0257-1
- [19] Kruse T, Neuschulz J, Wilhelm L et al. Prenatal Diagnosis of Robin Sequence: Sensitivity, Specificity, and Clinical Relevance of an Index for Micrognathia. *Cleft Palate Craniofac J* 2021 ; 58: 1012—1019. doi:10.1177/1055665620972297
- [20] Baujat G, Faure C, Zaouche A et al. Oroesophageal motor disorders in Pierre Robin syndrome. *J Pediatr Gastroenterol Nutr* 2001; 32: 297-302. doi: 10.1097/00005176-200103000-00012
- [21] Hinrichsen K. The early development of morphology and patterns of the face in the human embryo. *Adv Anat Embryol Cell Biol* 1985; 98: 1-79. doi:10.1007/978-3-642-70754-4
- [22] Maarse W, Berge SJ, Pistorius L et al. Diagnostic accuracy of transabdominal ultrasound in detecting prenatal cleft lip and palate: a systematic review. *Ultrasound Obst Gyn* 2010; 35: 495—502. doi:10.1002/uog.7472
- [23] Maarse W, Pistorius LR, Van Eeten WK et al. Prenatal ultrasound screening for orofacial clefts. *Ultrasound Obstet Gynecol* 2011 ; 38: 434-439. doi:10.1002/uog.8895
- [24] Offerdal K, Jebens N, Syvertsen T et al. Prenatal ultrasound detection of facial clefts: a prospective study of 49,314 deliveries in a non-selected population in Norway. *Ultrasound Obstet Gynecol* 2008; 31 639-646. doi:10.1002/uog.5280
- [25] Wilhelm L, Borgers H. The 'equals sign': a novel marker in the diagnosis of fetal isolated cleft palate. *Ultrasound Obstet Gynecol* 2010; 36: 439-444. doi:10.1002/uog.7704
- [26] Lind K, Aubry MC, Belarbi N et al. Prenatal diagnosis of Pierre Robin Sequence: accuracy and ability to predict phenotype and functional severity. *Prenat Diagn* 2015; 35: 853-858. doi:10.1002/pd.4619
- [27] Wiechers C, Poets C, Hoopmann M et al. Fetal Profile Markers for the Detection of Robin Sequence in Fetuses With Retrognathia. *Ultraschall Med* 2021. doi:10.1055/a-1659-2499
- [28] Bronshtein M, Blazer S, Zalel Y et al. Ultrasonographic diagnosis of glossoptosis in fetuses With Pierre Robin sequence in early and mid pregnancy. *Am J Obstet Gynecol* 2005; 193: 1561—1564. doi:10.1016/j.ajog.2005.03.052

- [29] Ball JV, DiBiase DD, Sommerlad BC. Transverse maxillary arch changes with the use of preoperative orthopedics in unilateral cleft palate infants. *Cleft Palate Craniofac J* 1995; 32: 483—488. doi:10.1597/1545-1569\_1995\_032\_0483\_tmacw\_2.3.co\_2
- [30] Winters JC, Hurwitz DJ. Presurgical orthopedics in the surgical management of unilateral cleft lip and palate. *Plast Reconstr Surg* 1995; 95: 755-764. doi:10.1097/00006534-199504000-00024
- [31] Uzel A, Alparslan ZN. Long-term effects of presurgical infant orthopedics in patients with cleft lip and palate: a systematic review. *Cleft Palate Craniofac J* 2011; 48: 587-595. doi:10.1597/10-008
- [32] Mishima K, Sugahara T, Mori Y et al. Three-dimensional comparison between the palatal forms in complete unilateral cleft lip and palate with and without Hotz plate from cheiloplasty to palatoplasty. *Cleft Palate Craniofac J* 1996; 33: 312-317. doi:10.1597/1545-1569\_1996\_033\_0312\_tdcctp\_2.3.co\_2
- [33] Hotz M, Gnoinski W. Comprehensive care of cleft lip and palate children at Zurich university: a preliminary report. *Am J Orthod* 1976; 70: 481-504. doi:10.1016/0002-9416(76)90274-8
- [34] Weise C, Frank K, Wiechers C et al. Intraoral scanning of neonates and infants with craniofacial disorders: feasibility, scanning duration, and clinical experience. *European Journal of Orthodontics* 2021 ; 1—8. doi:10.1093/ejo/cjab068
- [35] Sillman J. Dimensional changes of the dental arches: longitudinal Study from birth to 25 years. *American Journal of Orthodontics* 1964; 50: 824-842
- [36] Stahl F, Grabowski R, Gaebel M et al. Relationship between occlusal findings and orofacial myofunctional status in primary and mixed dentition. Part II: Prevalence of orofacial dysfunctions. *J Orofac Orthop* 2007; 68: 74-90. doi:10.1007/s00056-007-2606-9
- [37] Maull DJ, Grayson BH, Cutting CB et al. Long-term effects of nasoalveolar molding on three-dimensional nasal Shape in unilateral clefts. *Cleft Palate J* 1999; 36: 391-397. doi:10.1597/1545-1569\_1999\_036\_0391\_lteonm\_2.3.co\_2
- [38] Cohen MM Jr.. Robin sequences and complexes: causal heterogeneity and pathogeneticphenotypic variability. *Am J Med Genet* 1999; 84: 311-315
- [39] Poets CF, Maas C, Buchenau W et al. Multicenter Study on the effectiveness of the pre-epiglottic baton plate for airway obstruction and feeding problems in Robin sequence. *Orphanet Journal of Rare Diseases* 2017; 12: 1-6. doi:10.1186/s13023-017-0602-8
- [40] Bacher M, Sautermeister J, Urschitz MS et al. An oral appliance with velar extension for treatment of obstructive sleep apnea in infants with Pierre Robin sequence. *Cleft Palate Craniofac J* 2011 ; 48: 331-336. doi:10.1597/09-091
- [41] Nicholson P, Plint D. A long-term Study of rapid maxillary expansion and bone grafting in cleft lip and palate patients. *The European Journal of Orthodontics* 1989; 11: 186—192

- [42] Pucciarelli MGR, De Lima Toyoshima GH, Oliveira TM et al. Assessment of dental arch stability after orthodontic treatment and oral rehabilitation in complete unilateral cleft lip and palate and non-clefts patients using 3D stereophotogrammetry. *BMC Oral Health* 2020; 20: 1-8
- [43] Meazzini MC, Corradi F, Mazzoleni F et al. Circummaxillary Sutures in Patients With Apert, Crouzon, and Pfeiffer Syndromes Compared to Nonsyndromic Children: Growth, Orthodontic, and Surgical Implications. *Cleft Palate Craniofac J* 2021 ; 58: 299—305. doi: 10.1177/1055665620947616
- [44] Liou EJ-W, Tsai W-C. A new protocol for maxillary protraction in cleft patients: repetitive weekly protocol of alternate rapid maxillary expansions and constrictions. *The Cleft palate-craniofacial journal* 2005; 42: 121-127. doi:10.1597/03-107.1
- [45] Meazzini MC, Zappia LB, Tortora Cet al. Short-and long-term effects of late maxillary advancement With the Liou-Alt-RAMEC protocol in unilateral cleft lip and palate. *The Cleft Palate-Craniofacial Journal* 2019; 56: 159-167. doi:10.1177/1055665618772395
- [46] Da Silva Filho OG, Teles SG, Ozawa TO et al. Secondary bone graft and eruption of the permanent canine in patients With alveolar clefts: literature review and case report. *The Angle Orthodontist* 2000; 70: 174-178
- [47] Tanimoto K, Yanagida T, Tanne K. Orthodontic treatment with tooth transplantation for patients with cleft lip and palate. *The Cleft palate-craniofacial journal* 2010; 47: 499—506
- [48] Isono H, Kaida K, Hamada Y et al. The reconstruction of bilateral clefts using endosseous implants after bone grafting. *American journal of orthodontics and dentofacial orthopedics* 2002; 121 : 403—410
- [49] Oosterkamp BC, Dijkstra PIJ, Remmelink HJ et al. Orthodontic space closure versus prosthetic replacement of missing upper lateral incisors in patients With bilateral cleft lip and palate. *The Cleft palate- craniofacial journal* 2010; 47: 591 —596
- [50] Al-Tarawneh A, Alrbata R, Alazmi K et al. Orthodontic management of craniofacial anomalies, a summary of the available options. *International Journal of Dentistry and Oral Health* 2016; 2: 1
- [51] Posnick JC, Taylor M. Skeletal stability and relapse patterns after Le Fort I osteotomy using miniplate fixation in patients with isolated cleft palate. *Plast Reconstr Surg* 1994; 94: 51—58. discussion 59—60. doi: 10.1097/00006534-199407000-00004
- [52] Matsuda A, Suda N, Motohashi N et al. Skeletal characteristics and treatment outcome of five patients with Robin sequence. *Angle Orthod* 2006; 76: 898-908. doi:10.1043/0003-3219(2006)076[0898:SCATOO]2.0.CO;2
- [53] Suri S, Ross RB, Tompson BD. Craniofacial morphology and adolescent facial growth in Pierre Robin sequence. *Am J Orthod Dentofacial Orthop* 2010; 137: 763-774. doi:10.1016/j.ajodo.2008.07.020
- [54] Wiechers C, Buchenau W, Arand J et al. Mandibular growth in infants with Robin sequence treated With the Tübingen palatal Plate. *Head & Face Medicine* 2019; 15: 1-8



- [55] Pfaff MJ, De Leon F, Le L et al. Long-term orthognathic considerations in the Pierre Robin sequence patient. Plastic and Reconstructive Surgery 2020; 146: 599e-606e